



IHC image of enteroendocrine cells in the intestinal epithelium.

## Glucagon-like Protein Receptor (GLP2R) Antibody

<b>Catalog #</b>	24200	<b>Product type</b>	Primary antibodies
<b>Lot #</b>	1539001L	<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid (100 µL)	<b>Isotype</b>	IgG
<b>Host</b>	Rabbit	<b>Preservative</b>	≤ 0.05% sodium azide
<b>Reacts With</b>	Rat	<b>Antigen</b>	Glucagon-like peptide 2 receptor (GLP2R) for acetyl 65-88 amide sequence targeting rat and human proteins, but not mouse. The peptide was synthesized and cross-linked to keyhole limpet hemocyanin via sulfolink coupling.

### INSTRUCTIONS

<b>Preparation</b>	The antiserum is provided as 100 µL of affinity purified serum in PBS (0.02M sodium phosphate with 0.15 M sodium chloride, pH 7.5) with 1% BSA. Reconstitution is not required. Recommend briefly spinning tube (30 sec. 200xg) to collect contents at bottom of tube.  Refer to the Instruction Manual available online at <a href="http://www.immunostar.com">www.immunostar.com</a> for information on tissue preparation, immunostaining techniques, troubleshooting, and formulas.
--------------------	---

### APPLICATION

<b>Quality Control</b>	The ImmunoStar GLP2 receptor antibody was quality control tested using standard immunohistochemical methods. The antiserum demonstrates significant labeling of enteroendocrine cells in the intestinal epithelium, as well as cell bodies of vagal afferents in nodose ganglia of the parasympathetic nervous system. Labeling is effective using indirect immunofluorescent and biotin/avidin-HRP techniques. The addition of intensifying reagents such as tyramide amplification or nickel ammonium sulfate to the chromogen solution will approximately double the dilution factor as recommended. Immunolabeling of western blot revealed a band of approximately 66 kDa in human and rat tissue, but not mouse.
<b>Tissue</b>	Rat and Human intestinal epithelium and nodose ganglia.
<b>Perfusion Fixation</b>	<ul style="list-style-type: none"> <li>Fixative: 4% paraformaldehyde in 0.1M phosphate buffer, pH 7.4; 500 mL over 20 min.</li> <li>Post Fixation: 1.5 hour at 4°C in 4% paraformaldehyde in 0.1M phosphate buffer, pH 7.4.</li> </ul>
<b>Sections</b>	50 µm vibratome
<b>Tissue Incubation</b>	48 hours at 2°–8° C
<b>Detection System</b>	Use Bn/Av-HRP at dilutions recommended by the manufacturer.
<b>Suggested Dilution</b>	1/800-1/1000 in PBS/0.3% Triton X-100 – Bn/Av-HRP immunohistochemistry

### NOTES

<b>Special Instructions</b>	It is recommended that the researcher perform a primary antibody dilution series using our dilution recommendations as a guideline. Note that a change in the fixation or buffering system from our protocol may change the configuration of the protein which could alter the reactivity with the tissue tested.
<b>Storage</b>	Store at 2 °–8°C until expiration date.
<b>Concentration</b>	300 µg/ml
<b>Journal References</b>	<a href="http://www.immunostar.com/publications">www.immunostar.com/publications</a>

*For Laboratory Reagent Use Only. Analytical and performance characteristics are not established.*

ALL PRODUCTS ARE FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE RRID:AB\_572211