

## nNOS:N-Terminal Peptide Control

<b>Catalog #</b>	<b>24447</b>
<b>Lot #</b>	<b>926604B</b>
<b>Form</b>	Lyophilized
<b>Product Type</b>	Peptide

### INFORMATION

<b>IHC Quality Control</b>	<p>The peptide control for nNOS (N-terminal) is intended for the immunoabsorption of nNOS (N-terminal) antiserum, catalog number 24431. Preadsorption of nNOS (N-terminal) antiserum, diluted according to the antibody specification sheet, with 5 µg/ml nNOS (N-terminal) peptide immunogen following the instructions below provides complete blockage of nNOS (N-terminal) immunolabeling. The peptide is provided as 25 µg of lyophilized human nNOS (N-terminal), and ≤ 0.09% sodium azide, sequence (134-148).</p> <p>Refer to the Instruction Manual available online at <a href="http://www.immunostar.com">www.immunostar.com</a> for information on tissue preparation, immunostaining techniques, troubleshooting, and formulas.</p>
<b>Storage/Handling</b>	<ul style="list-style-type: none"> <li>• The product does not need to be kept cooled during shipping. Prior to reconstitution, as described below, store long-term at -15°C or lower. The crystals in the vial may be difficult to see and the vial may appear empty.</li> <li>• After reconstitution, use immediately or refrigerate at 2°–8°C up to 2 days. For long-term storage, aliquot reconstituted material as desired and freeze at -15°C or lower. Avoid repeated freeze/thaw cycles.</li> </ul>

### APPLICATION

<b>Instructions for Use</b>	<ol style="list-style-type: none"> <li>1. Reconstitute vial with 500 µl of PBS/Triton X-100 (0.05M sodium phosphate, 0.15M sodium chloride, 0.3% Triton X-100, pH 7.4), put the stopper back on and vortex lightly until dissolved.</li> <li>2. Add 100 µl of the reconstituted peptide per milliliter of diluted antibody to achieve a final peptide concentration of 5 µg/ml.</li> <li>3. Vortex the diluted antibody/peptide solution and incubate at 2°–8°C for 18–24 hours before use.</li> <li>4. Always run preadsorbed antibody in parallel with untreated antibody. Untreated antibody should stain appropriately, while preadsorbed antibody should provide complete blockage of immunolabeling.</li> </ol>

### NOTES

<b>Journal References</b>	<a href="http://www.immunostar.com/publications">www.immunostar.com/publications</a>
---------------------------	--

*For Laboratory Reagent Use Only. Analytical and performance characteristics are not established.*

**ALL PRODUCTS ARE FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE**