



IHC image of neurons in rat supraoptic nucleus.

C-FOS Antibody

Catalog #	26209	Product type	Primary antibodies
Lot #	113018B	Clonality	Polyclonal
Form	Lyophilized whole serum (100 µL)	Isotype	IgG
Host	Rabbit	Preservative	≤ 0.09% sodium azide
Reacts With	Mouse, Rat	Antigen	Synthetic peptide sequence corresponding to (human) C-FOS (4–17) coupled to BTg with glutaraldehyde

INSTRUCTIONS

Preparation	<p>Do not reconstitute until ready to use since the product is most stable when lyophilized. The product does not need to be kept cooled during shipping; however, for long-term storage, store lyophilized antibody until ready to use at -15°C or lower. Reconstitute with 100 µL of distilled or deionized water. After reconstitution, use immediately or refrigerate at 2°-8°C. To avoid freeze/thaw cycles, dilute unused antibody with PBS or Tris buffer at a dilution no higher than 1/10, then aliquot and freeze at -15°C or lower.</p> <p>Refer to the Instruction Manual available online at www.immunostar.com for information on tissue preparation, immunostaining techniques, troubleshooting, and formulas.</p>
--------------------	---

APPLICATION

IHC Quality Control	For induction of c-Fos protein activity, rats were injected with 1.0 ml of 1.5 M NaCl per 100 grams of body weight. Negative control rats were injected with the same volume of normal saline. The ImmunoStar c-Fos antiserum was quality control tested using standard immunohistochemical methods. The antiserum demonstrates significant labeling of rat paraventricular nucleus and supraoptic nucleus using indirect immunofluorescent and biotin/avidin-HRP techniques. No labeling was seen in negative control rats. Specificity of the antiserum was demonstrated by blockage of staining in experimental rats by omission of c-Fos antibody or by substitution of antibody pre-incubated with synthetic peptide or the conjugate.
Tissue	Rat paraventricular nucleus and supraoptic nucleus
Perfusion Fixation	<ul style="list-style-type: none"> Fixative: 4% paraformaldehyde in 0.1M Phosphate buffer, pH 7.4; 500 mL over 20–30 min. Post Fixation: 1.5 hour at 4°C in 4% paraformaldehyde in 0.1M phosphate buffer, pH 7.4.
Sections	50 µm vibratome
Tissue Incubation	18–24 hours at 2°–8°C.
Detection System	Use IF or Bn/Av-HRP reagents at dilutions recommended by the manufacturer.
Suggested Dilution	1/4,000–1/6,000 in PBS/0.3% Triton X-100 – Bn/Av-HRP immunohistochemistry

NOTES

Special Instructions	It is recommended that the researcher perform a primary antibody dilution series using our dilution recommendations as a guideline. Note that a change in the fixation or buffering system from our protocol may change the configuration of the protein which could alter the reactivity with the tissue tested.
Long-Term Storage	After reconstitution, use immediately or refrigerate at 2°–8°C up to 2 days. For long-term storage, aliquot and freeze at -15°C or lower. Avoid repeated freeze/thaw cycles.
Concentration	Not applicable. Antibody concentration is only relevant for purified antibodies.
Journal References	www.immunostar.com/publications

For Laboratory Reagent Use Only. Analytical and performance characteristics are not established.

ALL PRODUCTS ARE FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE

RRID:AB_572267